

# JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Before-and-After Photos (Brenda I)



**Actual patient. Results may vary.** Unretouched photos taken before treatment and 1 month after treatment. A total of 1.4 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area. A total of 2.9 mL of JUVÉDERM VOLLURE™ XC was injected into the parentheses and marionette lines. A total of 2.2 mL of JUVÉDERM VOLBELLA® XC was injected into the vertical lip lines and lips.

## APPROVED USES

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over 21.

JUVÉDERM VOLLURE™ XC injectable gel is for injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds, in adults over 21.

JUVÉDERM VOLBELLA® XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over 21.

## IMPORTANT SAFETY INFORMATION

### Are there any reasons why I should not receive any JUVÉDERM® injectable gel formulation?

Do not use these products if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the Gram-positive bacterial proteins used in these products.

### What precautions should my doctor advise me about?

- Tell your doctor if you are pregnant or breastfeeding. The safety of these products for use during pregnancy or while breastfeeding has not been studied
- The safety of JUVÉDERM VOLUMA® XC in patients under 35 years or over 65 years, and the safety of JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years, has not been studied
- The safety and effectiveness of JUVÉDERM VOLUMA® XC in areas other than the cheek area, JUVÉDERM VOLLURE™ XC for areas other than facial wrinkles and folds, and JUVÉDERM VOLBELLA® XC in areas other than the lips and perioral area have not been established in clinical studies
- Tell your doctor if you have a history of excessive scarring (eg, hypertrophic scarring and keloid formations) or pigmentation disorders, as use of these products may result in additional scars or changes in pigmentation
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment

- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on immunosuppressive therapy used to decrease the body's immune response, as use of these products may result in an increased risk of infection
- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners, as this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment

### What are possible side effects?

The most commonly reported side effects with JUVÉDERM® injectable gels included injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, most side effects were moderate and lasted 2 to 4 weeks. For JUVÉDERM VOLLURE™ XC, most side effects were mild or moderate and lasted 14 days or less. For JUVÉDERM VOLBELLA® XC, most side effects were mild or moderate and lasted 30 days or less.

One of the risks with using these products is unintentional injection into a blood vessel, and, while rare, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring.

As with all skin injection procedures, there is a risk of infection.

**To report a side effect with any JUVÉDERM® product, please call Allergan at 1-800-433-8871. Please visit [Juvederm.com](http://Juvederm.com) or talk to your doctor for more information.**

Available by prescription only.



© 2017 Allergan. All rights reserved.  
All trademarks are the property of their respective owners.  
[Juvederm.com](http://Juvederm.com) JUV105857 03/17 170327

# JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Before-and-After Photos (Brenda 2)



**Actual patient. Results may vary.** Unretouched photos taken before treatment and 1 month after treatment. A total of 1.4 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area. A total of 2.9 mL of JUVÉDERM VOLLURE™ XC was injected into the parentheses and marionette lines. A total of 2.2 mL of JUVÉDERM VOLBELLA® XC was injected into the vertical lip lines and lips.

## APPROVED USES

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over 21.

JUVÉDERM VOLLURE™ XC injectable gel is for injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds, in adults over 21.

JUVÉDERM VOLBELLA® XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over 21.

## IMPORTANT SAFETY INFORMATION

### Are there any reasons why I should not receive any JUVÉDERM® injectable gel formulation?

Do not use these products if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the Gram-positive bacterial proteins used in these products.

### What precautions should my doctor advise me about?

- Tell your doctor if you are pregnant or breastfeeding. The safety of these products for use during pregnancy or while breastfeeding has not been studied
- The safety of JUVÉDERM VOLUMA® XC in patients under 35 years or over 65 years, and the safety of JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years, has not been studied
- The safety and effectiveness of JUVÉDERM VOLUMA® XC in areas other than the cheek area, JUVÉDERM VOLLURE™ XC for areas other than facial wrinkles and folds, and JUVÉDERM VOLBELLA® XC in areas other than the lips and perioral area have not been established in clinical studies
- Tell your doctor if you have a history of excessive scarring (eg, hypertrophic scarring and keloid formations) or pigmentation disorders, as use of these products may result in additional scars or changes in pigmentation
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment

- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on immunosuppressive therapy used to decrease the body's immune response, as use of these products may result in an increased risk of infection
- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners, as this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment

### What are possible side effects?

The most commonly reported side effects with JUVÉDERM® injectable gels included injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, most side effects were moderate and lasted 2 to 4 weeks. For JUVÉDERM VOLLURE™ XC, most side effects were mild or moderate and lasted 14 days or less. For JUVÉDERM VOLBELLA® XC, most side effects were mild or moderate and lasted 30 days or less.

One of the risks with using these products is unintentional injection into a blood vessel, and, while rare, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring.

As with all skin injection procedures, there is a risk of infection.

**To report a side effect with any JUVÉDERM® product, please call Allergan at 1-800-433-8871. Please visit [Juvederm.com](http://Juvederm.com) or talk to your doctor for more information.**

Available by prescription only.



© 2017 Allergan. All rights reserved.  
All trademarks are the property of their respective owners.  
[Juvederm.com](http://Juvederm.com) JUV105857 03/17 170327

# JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Before-and-After Photos (Brenda 3)



**Actual patient. Results may vary.** Unretouched photos taken before treatment and 1 month after treatment. A total of 1.4 mL of JUVÉDERM VOLUMA® XC was injected into the cheek area. A total of 2.9 mL of JUVÉDERM VOLLURE™ XC was injected into the parentheses and marionette lines. A total of 2.2 mL of JUVÉDERM VOLBELLA® XC was injected into the vertical lip lines and lips.

## APPROVED USES

JUVÉDERM VOLUMA® XC injectable gel is for deep injection in the cheek area to correct age-related volume loss in adults over 21.

JUVÉDERM VOLLURE™ XC injectable gel is for injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds, in adults over 21.

JUVÉDERM VOLBELLA® XC injectable gel is for injection into the lips for lip augmentation and for correction of perioral lines in adults over 21.

## IMPORTANT SAFETY INFORMATION

### Are there any reasons why I should not receive any JUVÉDERM® injectable gel formulation?

Do not use these products if you have a history of multiple severe allergies or severe allergic reactions (anaphylaxis), or if you are allergic to lidocaine or the Gram-positive bacterial proteins used in these products.

### What precautions should my doctor advise me about?

- Tell your doctor if you are pregnant or breastfeeding. The safety of these products for use during pregnancy or while breastfeeding has not been studied
- The safety of JUVÉDERM VOLUMA® XC in patients under 35 years or over 65 years, and the safety of JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years, has not been studied
- The safety and effectiveness of JUVÉDERM VOLUMA® XC in areas other than the cheek area, JUVÉDERM VOLLURE™ XC for areas other than facial wrinkles and folds, and JUVÉDERM VOLBELLA® XC in areas other than the lips and perioral area have not been established in clinical studies
- Tell your doctor if you have a history of excessive scarring (eg, hypertrophic scarring and keloid formations) or pigmentation disorders, as use of these products may result in additional scars or changes in pigmentation
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment

- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on immunosuppressive therapy used to decrease the body's immune response, as use of these products may result in an increased risk of infection
- Tell your doctor if you are using medications that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners, as this may result in increased bruising or bleeding at the injection site
- Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment

### What are possible side effects?

The most commonly reported side effects with JUVÉDERM® injectable gels included injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, most side effects were moderate and lasted 2 to 4 weeks. For JUVÉDERM VOLLURE™ XC, most side effects were mild or moderate and lasted 14 days or less. For JUVÉDERM VOLBELLA® XC, most side effects were mild or moderate and lasted 30 days or less.

One of the risks with using these products is unintentional injection into a blood vessel, and, while rare, the complications can be serious and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring.

As with all skin injection procedures, there is a risk of infection.

**To report a side effect with any JUVÉDERM® product, please call Allergan at 1-800-433-8871. Please visit [Juvederm.com](http://Juvederm.com) or talk to your doctor for more information.**

Available by prescription only.



© 2017 Allergan. All rights reserved.

All trademarks are the property of their respective owners.

[Juvederm.com](http://Juvederm.com) JUV105857 03/17 170327

# JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Indications and Important Safety Information

## JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Important Information

### INDICATIONS

JUVÉDERM VOLUMA® XC injectable gel is indicated for deep (subcutaneous and/or supraperiosteal) injection for cheek augmentation to correct age-related volume deficit in the mid-face in adults over the age of 21.

JUVÉDERM VOLLURE™ XC injectable gel is indicated for injection into the mid-to-deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds) in adults over the age of 21.

JUVÉDERM VOLBELLA® XC injectable gel is indicated for injection into the lips for lip augmentation and for correction of perioral rhytids in adults over the age of 21.

### IMPORTANT SAFETY INFORMATION

#### CONTRAINDICATIONS

These products should not be used in patients who have severe allergies, marked by a history of anaphylaxis or history or presence of multiple severe allergies, and should not be used in patients with a history of allergies to Gram-positive bacterial proteins or lidocaine contained in these products.

#### WARNINGS

- Do not inject into blood vessels. Introduction of these products into the vasculature may lead to embolization, occlusion of the vessels, ischemia, or infarction. Take extra care when injecting soft-tissue fillers; for example, inject the product slowly and apply the least amount of pressure necessary. Rare, but serious, adverse events associated with the intravascular injection of soft-tissue fillers in the face have been reported and include temporary or permanent vision impairment, blindness, cerebral ischemia or cerebral hemorrhage leading to stroke, skin necrosis, and damage to underlying facial structures. Immediately stop the injection if a patient exhibits any of the following symptoms: changes in vision, signs of a stroke, blanching of the skin, unusual pain during or shortly after the procedure. Patients should receive prompt medical attention and, possibly, evaluation by an appropriate healthcare professional specialist should an intravascular injection occur
- Product use at specific sites in which an active inflammatory process (skin eruptions such as cysts, pimples, rashes, or hives) or infection is present should be deferred until the underlying process has been controlled

#### PRECAUTIONS

- In order to minimize the risk of potential complications, these products should only be used by healthcare professionals who have appropriate training, experience, and knowledge of facial anatomy
- Healthcare professionals are encouraged to discuss the potential risks of soft-tissue injections with their patients prior to treatment and ensure that patients are aware of signs and symptoms of potential complications

- The safety and effectiveness for the treatment of anatomic regions other than the mid-face with JUVÉDERM VOLUMA® XC, facial wrinkles and folds with JUVÉDERM VOLLURE™ XC, and the lips and perioral area with JUVÉDERM VOLBELLA® XC have not been established in controlled clinical studies
- As with all transcutaneous procedures, dermal filler implantation carries a risk of infection. Follow standard precautions associated with injectable materials
- The safety for use during pregnancy, in breastfeeding females, and in patients with known susceptibility to keloid formation, hypertrophic scarring, and pigmentation disorders has not been studied
- The safety for use of JUVÉDERM VOLUMA® XC in patients under 35 or over 65 years, and JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years has not been established
- Use with caution in patients on immunosuppressive therapy
- Patients who are using products that can prolong bleeding (such as aspirin, nonsteroidal anti-inflammatory drugs, and warfarin) may experience increased bruising or bleeding at treatment sites
- If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment, or if these products are administered before the skin has healed completely, there is a possible risk of an inflammatory reaction at the treatment site
- Patients who experience skin injury near the site of implantation may be at a higher risk for adverse events
- The safety of JUVÉDERM VOLUMA® XC injectable gel for use in patients with very thin skin in the mid-face has not been established
- Patients may experience late onset nodules with use of dermal fillers, including JUVÉDERM VOLUMA® XC
- Patients may experience late onset adverse events with use of dermal fillers

#### ADVERSE EVENTS

The most commonly reported side effects for JUVÉDERM® injectable gels were injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, side effects were predominantly moderate in severity, with a duration of 2 to 4 weeks; for JUVÉDERM VOLLURE™ XC, they were mostly mild or moderate in severity, with a duration of 14 days or less; and for JUVÉDERM VOLBELLA® XC, they were predominantly mild or moderate, with duration of 30 days or less.

**To report an adverse reaction with any product in the JUVÉDERM® Collection, please call Allergan at 1-800-433-8871. Please visit [JuvedermDFU.com](http://JuvedermDFU.com) for more information.**

JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC injectable gels are available by prescription only.

